Northwestern Medicine[®]



TESTS AND PROCEDURES

Cardiopulmonary Exercise Stress Test Arterial Line Insertion: After-Hospital Care

The following information will help you recover after your cardiopulmonary exercise stress test with an arterial line insertion.

If you have any questions or concerns about your test, please ask your physician or nurse.

General information

Insertion site care

- After the procedure, we will apply an Elastoplast® bandage made of stretching elastic material to the wound site. Keep that on for at least 1 hour after you have been discharged from the hospital. This is important because bleeding or swelling can occur if you bend your wrist.
- After you remove the bandage, you may leave the area open to air or apply a Band-Aid[®].
- Apply ice to the insertion site once you are home as needed. This will help reduce any swelling and decrease any discomfort you may have.
- Keep the site clean and dry.
- Avoid exposure to products such as cleaning solvents for the first 24 hours. They may irritate the wound.

Insertion site healing

- The wound should remain soft and dry. A bruise (black and blue) or a small, marble-sized lump may be present.
- Please notify your physician if you have any of the following symptoms:
 - Redness around the puncture site
 - Drainage from the puncture site
 - A lump at the puncture site that enlarges or is larger than a marble
 - Pain at the puncture site that makes bending your wrist difficult
 - Discoloration or coolness of your hand

Warning signs and symptoms

If you notice minor bleeding from the puncture site, please do the following:

- Apply and hold firm pressure on the site for 10 minutes. You may use a clean cloth or tissue to apply pressure. After 10 minutes, remove pressure. The wound should be dry and flat, without bleeding.
- Cover the wound with a Band-Aid.
- Go to the nearest Emergency Department if you have any of these symptoms in your affected arm:
 - Bleeding that does not stop after you apply pressure
 - Swelling or change in color
 - Pain, numbness or tingling

If a problem occurs, or if you have questions, please call the clinical nurse specialist at 312.926.1583 (TTY: 711).