MWLC EMS Skill Performance Record OROPHARYNGEAL SUCTIONING

Name:	1 st attempt:	Pass	Repeat
Date:	2 nd attempt:	Pass	Repeat

Instructions: An adult's mouth is filled with blood. You are asked to assemble the equipment, choose the correct catheter from those available, and perform oropharyngeal suctioning.

Equipment needed: Airway manikin; various sizes suction catheters, suction unit, BSI

	quipment needed. Aliway mankin, various sizes suction catheters, suction dritt, Bot				
	Performance standard O Step omitted (or leave blank) 1 Not yet competent: Unsuccessful; required critical or excess prompting; marginal or inconsistent technique. 2 Successful; competent with correct timing, sequence & technique, no prompting necessary.	Attempt 1 rating	Attempt 2 rating		
	State indications for procedure: Secretions in mouth, nose or pharynx				
	* Universal plus droplet precautions (gloves/face shield)				
	Prepare patient Explain steps of procedure to patient				
	Obtain SpO₂ on room air if available and time allows				
	* Preoxygenate patient prior to suctioning if time allows				
	Prepare equipment: Inspect suction unit for power and proper assemblage				
	* Select appropriate suction catheter (flexible or rigid); attach to suction tubing				
	Perform procedure Open mouth using cross-finger technique				
	☐ Turn power on to high.☐ Kink tubing and ensure that unit achieves vacuum of 300 mmHg.				
	Without applying suction ☐ Insert suction catheter no deeper than pharynx. ☐ If Yankauer tip, insert w/ convex side along roof of mouth.				
	* Apply suction using a gentle twisting motion while limiting suction application to 10 sec on an adult and 5 sec in a child				
ľ	Refrain from jabbing catheter up and down while applying suction				
	* Reoxygenate patient with O₂ 15 L/NRM or BVM				
	Verbalize: Flush the suction catheter with NS or water between suction attempts to remove any material that could clog ports		A-8-000 - 8-00 - 1		
	Verbalize 2 complications if suction were applied improperly or for too long: □ *Hypoxia □ Atelectasis □ *Bradycardia □ Hypotension □ Tissue trauma □ ↑ ICP				
	Critical Criteria: Check if occurred during an attempt ☐ Failure to take or verbalize appropriate body substance isolation precautions ☐ Contaminates equipment or site without appropriately correcting the situation ☐ Performs any improper technique resulting in the potential for patient harm ☐ Exhibits unacceptable affect with patient or other personnel				
Sc	All steps must be independently performed in correct sequence with appropriate timing and all state explained/ performed correctly in order for the person to demonstrate competency. Any errors or or will require additional practice and a repeat assessment of skill proficiency.	arred (*) item missions of th	ns must be nese items		
	high quality without critical error, assistance or instruction.	, with exper	tise and to		
 Competent: Satisfactory performance without critical error; minimal coaching needed. Practice evolving/not yet competent: Did not perform in correct sequence, timing, and/or without prompts, relia manual, and/or critical error; recommend additional practice 					
		175 h 1 4 h 4 F			